

Extending Chacon-Walsh: Minimality and Generalised Starting Distributions

A. M. G. Cox*

Department of Mathematics,
University of York,
York YO10 5DD, U. K.

February 1, 2008

Abstract

In this paper¹ we consider the Skorokhod embedding problem for general starting and target measures. In particular, we provide necessary and sufficient conditions for a stopping time to be minimal in the sense of Monroe (1972). The resulting conditions have a nice interpretation in the graphical picture of Chacon and Walsh (1976).

Further, we demonstrate how the construction of Chacon and Walsh can be extended to any (integrable) starting and target distributions, allowing the constructions of Azema-Yor, Vallois and Jacka to be viewed in this context, and thus extended easily to general starting and target distributions. In particular, we describe in detail the extension of the Azema-Yor embedding in this context, and show that it retains its optimality property.

1 Introduction

The Skorokhod embedding problem has a long history, and was first posed (and solved) in Skorokhod (1965). Simply stated it is the following: given a stochastic process $(X_t)_{t \geq 0}$ and a distribution μ , find a stopping time T such that $X_T \sim \mu$.

In this work we will be interested in the case where B_t is a Brownian motion on $\mathbb{R}$, with a given (integrable) starting distribution μ_0 . Since Brownian motion on $\mathbb{R}$ is recurrent, the existence of such a stopping time is trivial: consider an independent random variable Y with distribution μ and run until the first time that the Brownian motion hits Y . Hence interest lies in the properties of the stopping time T and also of the stopped process $B_{t \wedge T}$.

*The author wishes to thank David Hobson for his helpful suggestions and advice. e-mail: amgc500@york.ac.uk; web: www-users.york.ac.uk/~amgc500/

¹MSC 2000 subject classifications. Primary: 60G40, 60J60; Secondary: 60G44, 60J65.

Keywords: Brownian Motion, Embedding, Azema-Yor Embedding, Stopping Time, Minimal Stopping Time, Chacon-Walsh Construction, Balayage.

Classically, the 1-dimensional question has been considered in the case where $B_0 = 0$, and the target distribution μ is centred. In this case many solutions are known: Azéma and Yor (1979); Bertoin and Le Jan (1992); Dubins (1968); Jacka (1988); Perkins (1986); Root (1969); Vallois (1983). We refer the reader to Oblój (2004) for an excellent recent survey of these results. A property shared by all of these embeddings is that the process $B_{t \wedge T}$ is uniformly integrable, and we shall call stopping times for which this is the case UI stopping times. Further, within the class of embeddings where T is UI, many of these stopping times have optimality properties: for example the Azema-Yor embedding maximises the law of the maximum, while the Vallois construction can be used to minimise or maximise $\mathbb{E}(f(L_T))$ for a convex function f (Vallois, 1992). It is clear that either of the maximisation problems are degenerate when looked at outside this class.

The class of UI stopping times can also be characterised in the following way due to Monroe (1972). We make the following definition:

Definition 1.1. A stopping time T for the process X is *minimal* if whenever $S \leq T$ is a stopping time such that X_S and X_T have the same distribution then $S = T$ a.s..

Then the class of minimal stopping times can be shown to be equivalent to the class of UI embeddings we had before:

Theorem 1.2. (Monroe, 1972, Theorem 3) *Let S be a stopping time such that $\mathbb{E}(B_S) = 0$. Then S is minimal if and only if the process $B_{t \wedge S}$ is uniformly integrable.*

Such a characterisation gives a natural interpretation to the class of UI embeddings.

Our interest in this paper lies in the extension to general starting measures. In such an example, even if the means agree, there is no guarantee that there will exist a UI stopping time which has the given starting and target distributions. This can be seen by considering the example of a target distribution consisting of a point mass at zero, but with starting distribution of mass $\frac{1}{2}$ at each of -1 and 1 . Clearly the only minimal stopping time is to stop the first time the process hits 0, however this stopping time is not UI.

In Cox and Hobson (2003) conditions for a stopping time to be minimal were considered. When the Brownian motion starts at the origin, and the target distribution is not centred, conditions on the process can be given which are equivalent to the stopping time being minimal. One of the main results of this work is to show that the conditions have an extension to the case of a general starting distribution, however the simple example given above shows that the extension is not trivial.

It will turn out that the characterisation of minimal stopping times is closely connected to the potentials of the two measures. In this context, the relationship between the measures can be viewed graphically in the framework of Chacon and Walsh (1976). In this paper a graphical construction is interpreted as a sequence of exit times from compact intervals, whose limit is an embedding. This is done for starting and target distributions which satisfy the relationship

$$-\mathbb{E}^{\mu_0}|X - x| = u_{\mu_0}(x) \geq u_{\mu}(x) = -\mathbb{E}^{\mu}|X - x| \quad (1)$$

for all $x \in \mathbb{R}$. We shall show that the construction can be extended to the case where this condition fails, and that the exact method of the extension will determine whether the stopping time is minimal.

Establishing this connection will then allow us to extend several existing embeddings (Azéma and Yor, 1979; Jacka, 1988; Vallois, 1992), to the more general setting (maintaining minimality) via a limiting argument.

2 The Balayage Construction

In the theory of general Markov processes, a common definition of the potential of a stochastic process is given by

$$U\mu(x) = \int_{\mathbb{R}} \mu(dy) \int_{\mathbb{R}^+} ds p_s(x, y),$$

where $p_s(x, \cdot)$ is the transition density at time s of the process started at x . In the case of Brownian motion, we note that the integral is infinite. To resolve this we use the compensated definition (and introduce new notation to emphasise the fact that this is not the classical definition of potential):

$$u_\mu(x) = \int_{\mathbb{R}} \mu(dy) \int_{\mathbb{R}^+} ds (p_s(x, y) - p_s(0, 0)).$$

This definition simplifies to the following:

$$u_\mu(x) = - \int |x - y| \mu(dy). \quad (2)$$

Remark 2.1. The function u_μ has the following properties:

- (i) The measure μ is integrable if and only if the function u_μ is finite for any (and therefore all) $x \in \mathbb{R}$.
- (ii) u_μ is continuous, differentiable everywhere except the set $\{x \in \mathbb{R} : \mu(\{x\}) > 0\}$ and concave.
- (iii) Write

$$m = \int x \mu(dx).$$

As $|x| \rightarrow \infty$, we have

$$u_\mu(x) + |x| \rightarrow m \operatorname{sign}(x). \quad (3)$$

- (iv) As a consequence of (3), if μ and ν are integrable distributions, then there exists a constant $K > 0$ such that:

$$\sup_{x \in \mathbb{R}} |u_\mu(x) - u_\nu(x)| < K.$$

(v) u_μ is almost everywhere differentiable with left and right derivatives

$$\begin{aligned} u'_{\mu,-}(x) &= 1 - 2\mu((-\infty, x)); \\ u'_{\mu,+}(x) &= 1 - 2\mu((-\infty, x]). \end{aligned}$$

Chacon (1977) contains many results concerning these potentials. We will describe a balayage technique that produces a sequence of measures and corresponding stopping times, and which will have as its limit our desired embedding. The following lemma will therefore be important in concluding that the limit we obtain will indeed be the desired distribution:

Lemma 2.2 (Chacon (1977), Lemmas 2.5, 2.6). *Suppose $\{\mu_n\}$ is a sequence of probability measures. If*

- (i) μ_n converges weakly to μ and $\lim_{n \rightarrow \infty} u_{\mu_n}(x_0)$ exists for some $x_0 \in \mathbb{R}$, then $\lim_{n \rightarrow \infty} u_{\mu_n}(x)$ exists for all $x \in \mathbb{R}$ and there exists $C \geq 0$ such that

$$\lim_{n \rightarrow \infty} u_{\mu_n}(x) = u_\mu(x) - C. \quad (4)$$

- (ii) $\lim_{n \rightarrow \infty} u_{\mu_n}(x)$ exists for all $x \in \mathbb{R}$ then μ_n converges weakly to μ for some measure μ and μ is uniquely determined by the limit $\lim_n u_{\mu_n}(x)$.

We consider the embedding problem where we have a Brownian motion B with $B_0 \sim \mu_0$ (an integrable starting distribution) and we wish to embed an integrable target distribution μ . This is essentially the case considered by Chacon and Walsh (1976), although they only consider the case where $u_{\mu_0}(x) \geq u_\mu(x)$ for all x (when (3) implies μ_0 and μ have the same mean) — we will see that this case is simpler than the general case we consider. The embedding problem is frequently considered when μ_0 is the Dirac measure at 0. One of the appealing properties of the case where $B_0 = 0$ is that for all centred target distributions (Chacon, 1977, Lemma 2.1)

$$u_\mu(x) \leq -|x| = u_{\mu_0}(x), \quad (5)$$

and the condition on the ordering of potentials is easily satisfied.

We extend the technique of Chacon and Walsh (1976) to allow balayage on semi-infinite intervals. This extra step in the construction allows further flexibility later when we take limits of the constructions. In particular it will make the application of subsequent results trivial. Each step in the construction is described mathematically by a simple balayage technique:

Definition 2.3. Let μ be a probability measure on $\mathbb{R}$, and I a finite, open interval, $I = (a, b)$. Then define the *balayage* μ_I of μ on I by:

$$\begin{aligned} \mu_I(A) &= \mu(A) & A \cap \bar{I} &= \emptyset; \\ \mu_I(\{a\}) &= \int_{\bar{I}} \frac{b-x}{b-a} \mu(dx); \\ \mu_I(\{b\}) &= \int_{\bar{I}} \frac{x-a}{b-a} \mu(dx); \\ \mu_I(I) &= 0. \end{aligned}$$

Suppose now $I = (a, \infty)$ (resp. $I = (-\infty, a)$), and define the *balayage* μ_I of μ by

$$\begin{aligned}\mu_I(A) &= \mu(A) & A \cap \bar{I} &= \emptyset; \\ \mu_I(\{a\}) &= \int_{\bar{I}} \mu(dx); \\ \mu_I(I) &= 0.\end{aligned}$$

The balayage μ_I is a probability measure and if I is a finite interval the means of μ and μ_I agree. In particular, μ_I is the law of a Brownian motion started with distribution μ and run until the first exit from I .

Our reason for introducing the Balayage technique is that the potential of μ_I is readily calculated from the potential of μ :

Lemma 2.4 (Chacon (1977) Lemma 8.1). *Let μ be a probability measure with finite potential, $I = (a, b)$ a finite open interval and μ_I the balayage of μ with respect to I . Then*

- (i) $u_\mu(x) \geq u_{\mu_I}(x) \quad x \in \mathbb{R};$
- (ii) $u_\mu(x) = u_{\mu_I}(x) \quad x \in I^c;$
- (iii) u_{μ_I} is linear for $x \in \bar{I}$.

When I is a semi-infinite interval we may calculate the potential in a similar way:

Lemma 2.5. *Let μ be a probability measure with finite potential u_μ , $I = (-\infty, a)$ or $I = (a, \infty)$ a semi-infinite interval and μ_I the balayage of μ with respect to I . Then*

$$\begin{aligned}u_{\mu_I}(x) &= u_\mu(x) + \Delta m & x &\notin I; \\ u_{\mu_I}(x) &= u_\mu(a) + \Delta m - |a - x| & x &\in I,\end{aligned}$$

where we have written

$$\Delta m = \int_I |x - a| \mu(dx).$$

The semi-infinite balayage step in Definition 2.3 can be recreated using the balayage steps on compact intervals, for example by taking the sequence of intervals $(a, a+1), (a, a+2), (a, a+3), \dots$. However it does not let us make the same constructions as we can with the extended definition — for example if we wish our first step to be to move up to 1, we would not be able to carry out any further steps.

Formally, we may use balayage to define an embedding as the following result shows. In the formulation of the result we assume we are given the sequence of functions we use to construct the stopping time, and from these deduce the target distribution. However we will typically use the result in situations where we have a desired target distribution and choose the sequence to fit this distribution.

Lemma 2.6. *Let $f_1, f_2, \dots$ be a sequence of linear functions on $\mathbb{R}$ such that $|f'_n(x)| \leq 1$ and*

$$g(x) = \inf_{n \in \mathbb{N}} f_n(x) \wedge (u_{\mu_0}(x)). \quad (6)$$

Set $T_0 = 0$, $g_0(x) = u_{\mu_0}(x)$ and, for $n \geq 1$, define

$$\begin{aligned} a_n &= \inf\{x \in \mathbb{R} : f_n(x) < g_{n-1}(x)\}; \\ b_n &= \sup\{x \in \mathbb{R} : f_n(x) < g_{n-1}(x)\}; \\ T_n &= \inf\{t \geq T_{n-1} : B_t \notin (a_n, b_n)\}; \\ g_n(x) &= g_{n-1}(x) \wedge f_n(x). \end{aligned}$$

Then the T_n are increasing so we define $T = \lim_{n \rightarrow \infty} T_n$. If

$$g(x) = u_\mu(x) - C \tag{7}$$

for some $C \in \mathbb{R}$ and some integrable probability measure μ then $T < \infty$ a.s. and T is an embedding of μ .

If we only consider the theorem under the condition $|f'_n(x)| < 1$ this is a formalised statement of the construction implicit in Chacon (1977) and made explicit under the further condition (1) in Chacon and Walsh (1976). The introduction of the balayage steps on the half-line is the novel content of the result.

Proof. The hard part is to show that if (7) holds then the stopping time T is almost surely finite. We prove in fact that $\mathbb{E}(L_T) < \infty$, where L is the local time of B at zero. By considering the martingale $|B_t| - L_t$ we must have

$$\mathbb{E}(L_{T_1}) = u_{\mu_0}(0) - f_1(0) \wedge u_{\mu_0}(0) \tag{8}$$

when the interval (a_1, b_1) is compact; by approximating the semi-infinite interval by compact intervals (and a monotone convergence argument) this will extend to all possible choices of f_1 , and, by an induction argument, we deduce

$$\mathbb{E}(L_{T_n}) = u_{\mu_0}(0) - \inf_{k \leq n} f_k(0) \wedge u_{\mu_0}(0).$$

A monotone convergence argument allows us to deduce that

$$\mathbb{E}(L_T) = u_{\mu_0}(0) - \inf_{n \in \mathbb{N}} f_n(0) \wedge u_{\mu_0}(0)$$

which is finite by (7), and hence $T < \infty$ a.s..

The functions g_n correspond to a potential of a measure μ_n (μ_n being the law of B_{T_n}) via:

$$g_n(x) = u_{\mu_n}(x) - C_n$$

for some constant C_n , and hence we have

$$u_{\mu_0}(x) \geq u_{\mu_n}(x) - C_n \geq u_\mu(x) - C$$

and as $n \rightarrow \infty$ the last two terms converge. From (3) we can deduce that

$$C_n \geq \left| \int x \mu_0(dx) - \int x \mu_n(dx) \right| \geq 0,$$

so that since $g_n(x) \geq g(x)$, we have

$$u_\mu(x) - C \leq u_{\mu_n}(x) \leq 0,$$

the second inequality coming from the definition of the potential. Consequently, taking $x = 0$, we can find a subsequence n_j for which $\lim_{j \rightarrow \infty} u_{\mu_{n_j}}(0)$ exists, and hence for which $\lim_{n \rightarrow \infty} C_{n_j}$ also exists. Since $g_n(x)$ converges pointwise to $g(x)$ we must also have pointwise convergence of $u_{\mu_{n_j}}(x)$ to $u_\mu(x) - C'$ for some constant C' , and by Lemma 2.2 $B_{T_{n_j}}$ converges weakly to μ . Since also $T_{n_j} \uparrow T$, by the continuity of the Brownian motion B_T has law μ . $\square$

The case considered by Chacon and Walsh (1976) has a notable property. When the starting and target measures are centred (or at least when their means agree) and

$$u_{\mu_0}(x) \geq u_\mu(x) \tag{9}$$

then we may choose a construction such that $C = 0$ in (7). In this case the process $B_{t \wedge T}$ is uniformly integrable (Chacon, 1977, Lemma 5.1). The desire to find a condition to replace uniform integrability in situations where (9) does not hold, and to construct suitable stopping times using this framework, is the motivation behind the subsequent work.

We note also that — for given μ, μ_0 — we may find a construction for any C which satisfies $C \geq \sup_x \{u_\mu(x) - u_{\mu_0}(x)\}$; as we shall see, the case where there is equality is of particular interest. As a consequence of (3) we must always have $C \geq 0$.

3 Minimality: Some Preliminary Results

In this and the subsequent section we discuss necessary and sufficient conditions for an embedding of an integrable target distribution to be minimal (Definition 1.1) when we have an integrable starting distribution. These results will extend the conditions of Theorems 1.2 and the following result:

Theorem 3.1 (Cox and Hobson (2003)). *Let T be a stopping time of Brownian motion which embeds an integrable distribution μ where $m = \int_{\mathbb{R}} x \mu(dx) < 0$. Then the following conditions are equivalent:*

(i) T is minimal for μ ;

(ii) for all stopping times $R \leq S \leq T$,

$$\mathbb{E}(B_S | \mathcal{F}_R) \leq B_R \quad a.s.; \tag{10}$$

In the case where $\text{supp}(\mu) \subseteq [\alpha, \infty)$ for some $\alpha < 0$ then the above conditions are also equivalent to the condition:

(iii)

$$\mathbb{P}(T \leq H_\alpha) = 1, \tag{11}$$

where $H_\alpha = \inf\{t \geq 0 : B_t = \alpha\}$ is the hitting time of α .

Remark 3.2. For further necessary and sufficient conditions, see also Cox and Hobson (2003).

As a starting point, we recall:

Proposition 3.3 (Monroe (1972), Proposition 2). *For any stopping time T there exists a minimal stopping time $S \leq T$ such that $B_S \sim B_T$.*

Monroe's proof does not rely on the fact that B starts at 0, and so the result extends to a general starting distribution.

It can also be seen that the argument used in Monroe (1972) to show that if the process is uniformly integrable then the process is minimal does not require the starting measure to be a point mass. For completeness we state a similar result, with the proof given in Monroe (1972):

Lemma 3.4. *Let T be a stopping time embedding μ in $(B_t)_{t \geq 0}$, with $B_0 \sim \mu_0$ where μ and μ_0 are integrable distributions. If*

$$\mathbb{E}(B_T | \mathcal{F}_S) = B_S \text{ a.s.} \quad (12)$$

for all stopping times $S \leq T$ then T is minimal.

Note that $S \equiv 0$ implies that μ, μ_0 have the same mean.

Remark 3.5. We will later be interested also in necessary conditions for minimality. The condition in (12) is not necessary even when both starting and target measures are centred, as can be seen by taking $\mu_0 = \frac{1}{2}\delta_{-1} + \frac{1}{2}\delta_1$ and $\mu = \delta_0$, where it is impossible to satisfy (12) but the (only) minimal stopping time is 'stop when the process hits 0.'

The condition in (12) is equivalent to uniform integrability of the process $(B_{t \wedge T})_{t \geq 0}$. One direction follows from the optional stopping theorem, the reverse implication comes from the upward martingale theorem (Rogers and Williams, 2000)[Theorem II.69.5], which tells us that the process $X_t = \mathbb{E}(B_T | \mathcal{F}_t)$ is a uniformly integrable martingale on $t \leq T$. When (12) holds, $X_t = B_{t \wedge T}$, and the process $B_{t \wedge T}$ is a uniformly integrable martingale.

For the rest of this section we will consider minimality for general starting and target measures: particularly when the means do not agree. If this occurs when the starting measure is a point mass, necessary and sufficient conditions are given in Theorem 3.1. In subsequent proofs with general starting measures we will often reduce problems to the point mass case in order to apply the result.

Remark 3.6. The condition given in (iii) of Theorem 3.1 hints at a more general idea inherent in the study of embeddings in Brownian motion. When $B_0 = 0$, it is a well known fact that if there exists $\alpha < 0 < \beta$ such that $T \leq H_\alpha \wedge H_\beta$ then $B_{t \wedge T}$ is a uniformly integrable martingale. If $T \leq H_\alpha$ then the process is a supermartingale. In terms of embeddings, this observation has the following consequence: if the target distribution is centred and supported on a bounded interval, an embedding is minimal if and only if the process never leaves this

interval. If the target distribution has a negative mean, but still lies on a bounded interval, any embedding must move above the interval — i.e. $\mathbb{P}(\sup_{t \leq T} B_t \geq x) > 0$ for all $x \geq 0$. Theorem 3.1 and Proposition 3.3 tell us that in this case an embedding exists for which $T \leq H_\alpha$ and all minimal embeddings satisfy this property.

Recall that there is a natural ordering on the set of (finite) measures on $\mathbb{R}$, that is $\mu \preceq \nu$ if and only if $\mu(A) \leq \nu(A)$ for all $A \in \mathcal{B}(\mathbb{R})$, in which case we say that ν dominates μ . In such instances it is possible to define a (positive, finite) measure $(\nu - \mu)(A) = \nu(A) - \mu(A)$. The notation $\nu = \mathcal{L}(B_T; T < H_\alpha)$ is used to mean the (sub-probability) measure ν such that $\nu(A) = \mathbb{P}(B_T \in A, T < H_\alpha)$.

Lemma 3.7. *Let B_t be a Brownian motion with $B_0 = 0$, T a stopping time embedding a distribution μ , $\tilde{\mu}$ a target distribution such that $\text{supp}(\tilde{\mu}) \subseteq [\alpha, \infty)$ for some $\alpha < 0$ and $\int x \tilde{\mu}(dx) \leq 0$. Then if $\nu = \mathcal{L}(B_T; T < H_\alpha)$ is dominated by $\tilde{\mu}$, there exists a minimal stopping time $\tilde{T} \leq T \wedge H_\alpha$ which embeds $\tilde{\mu}$.*

Similarly, if $\tilde{\mu}$ is such that $\text{supp}(\tilde{\mu}) \subseteq [\alpha, \beta]$ and $\int x \tilde{\mu}(dx) = 0$, and if $\nu = \mathcal{L}(B_T; T < H_\alpha \wedge H_\beta)$ is dominated by $\tilde{\mu}$, then there exists a minimal stopping time $\tilde{T} \leq T \wedge H_\alpha \wedge H_\beta$ which embeds $\tilde{\mu}$.

Proof. Construct a stopping time T' as follows: on $\{T < H_\alpha\}$, $T' = T$; otherwise choose T' so that $T' = H_\alpha + T'' \circ \theta_{H_\alpha}$ where T'' is chosen to embed $(\tilde{\mu} - \nu)$ on $\{T' \geq H_\alpha\}$ given $B_0 = \alpha$. Then T' is an embedding of $\tilde{\mu}$ and $T' \leq T$ on $\{T < H_\alpha\}$. So by Proposition 3.3 we may find a minimal embedding $\tilde{T} \leq T' \wedge H_\alpha = T \wedge H_\alpha$ which embeds $\tilde{\mu}$.

The proof in the centred case is essentially identical, but now stopping the first time the process leaves $[\alpha, \beta]$. $\square$

We turn now to the case of interest — that is when $B_0 \sim \mu_0$ and $B_T \sim \mu$ for integrable measures μ_0 and μ . The following lemma is essentially technical in nature, but will allow us to deduce the required behaviour on letting A increase in density.

Lemma 3.8. *Let T be a minimal stopping time, and A a countable subset of $\mathbb{R}$ such that A has finitely many elements in every compact subset of $\mathbb{R}$ and $d(x, A) < M$ for all $x \in \mathbb{R}$ and some $M > 0$. We consider the stopping time*

$$R(A) = \inf\{t \geq 0 : B_t \in A\} \wedge T$$

and we write

$$E_A(x) = \begin{cases} \mathbb{E}(B_T | T > R(A), B_{R(A)} = x) & : \mathbb{P}(T > R(A), B_{R(A)} = x) > 0; \\ x & : \mathbb{P}(T > R(A), B_{R(A)} = x) = 0. \end{cases}$$

Then there exists $a \in \bar{\mathbb{R}} = \mathbb{R} \cup \{-\infty\} \cup \{\infty\}$ such that

$$E_A(x) > x \implies x < a, \tag{13}$$

$$E_A(x) < x \implies x > a, \tag{14}$$

and $T \leq H_a$ on $\{T \geq R(A)\}$.

Further, if there exists $x < y$ such that $E_A(x) > x$ and $E_A(y) < y$ then there exists $a_\infty \in [x, y]$ such that $T \leq H_{a_\infty}$.

Proof. Suppose that there exists $x < y$ such that $E_A(x) < x$ and $E_A(y) > y$, and suppose $E_A(w) = w$ for $x < w < y$. We show that we can construct a strictly smaller embedding, contradicting the assumption that T is minimal.

Define the stopping time $T' = R(A)\mathbf{1}_{\{B_{R(A)} \in \{x, y\}\}} + T\mathbf{1}_{\{B_{R(A)} \notin \{x, y\}\}}$ and for some $z \in (x, y)$, the stopping time

$$T'' = \inf\{t \geq T' : B_t = z\} \wedge T.$$

As a consequence of Remark 3.6, paths from both x and y must hit z .

Consider the set $\{T'' < T\}$. On this set we have only paths with $B_{R(A)} = x$ and $B_{R(A)} = y$. Define $\mu_x = \mathcal{L}(B_T; B_{R(A)} = x, T'' < T)$ and $\mu_y = \mathcal{L}(B_T; B_{R(A)} = y, T'' < T)$. Since Brownian motion bounded above is a submartingale,

$$\mathbb{E}(B_{T \wedge H_z}; B_{R(A)} = x, T > R(A)) \geq x\mathbb{P}(B_{R(A)} = x, T > R(A)).$$

Together with $E_A(x) < x$ this implies

$$z\mathbb{P}(B_{R(A)} = x, T'' < T) > \mathbb{E}(B_T; B_{R(A)} = x, T'' < T),$$

i.e. we must have $\frac{1}{\mu_x(\mathbb{R})} \int w \mu_x(dw) < z$, and similarly $\frac{1}{\mu_y(\mathbb{R})} \int w \mu_y(dw) > z$. Then we apply Lemma 3.7 to the processes $B_{T''+t}$ on $\{B_{R(A)} = x, T'' < T\}$ and $\{B_{R(A)} = y, T'' < T\}$ with the measures

$$\begin{aligned} \tilde{\mu}_x &= \mu_x|_{[a_1, \infty)} + \mu_y|_{(a_2, \infty)} \\ \tilde{\mu}_y &= \mu_x|_{(-\infty, a_1)} + \mu_y|_{(-\infty, a_2)} \end{aligned}$$

where we choose $a_1 < z < a_2$ so that

$$\frac{1}{\tilde{\mu}_x(\mathbb{R})} \int w \tilde{\mu}_x(dw) \leq z \text{ and } \frac{1}{\tilde{\mu}_y(\mathbb{R})} \int w \tilde{\mu}_y(dw) \geq z$$

and also so that $\mu_x(\mathbb{R}) = \tilde{\mu}_x(\mathbb{R})$ and $\mu_y(\mathbb{R}) = \tilde{\mu}_y(\mathbb{R})$ ². This will produce a strictly smaller embedding, in contradiction to the assumption that T is minimal.

So we have shown that there exists a such that (13) and (14) hold. We just need to show that we can choose a so that $T \leq H_a$ on $\{T \geq R(A)\}$.

Suppose that there exists $x < y$ such that $E_A(x) > x$ and $E_A(y) < y$ and $E_A(w) = w$ for $w \in (x, y)$. If

$$\sup_{x < a} \mu_x((a, \infty)) = 0 \text{ and } \sup_{y > a} \mu_y((-\infty, a)) = 0 \text{ for some } a \in (x, y) \quad (15)$$

then T minimal and Theorem 3.1 implies that $T \leq H_a$ on $\{T \geq R(A)\}$.

²It may be necessary to consider only a proportion of the paths hitting z from one side; this can be done by choosing paths according to an independent $U([0, 1])$ random variable and running the rest of the paths according to T . This will still construct a strictly smaller stopping time.

So suppose that (15) does not hold. We shall show that we can then find a sequence $x_1, x_2, \dots, x_r$ of elements of A such that we are able to transfer mass between the x_i to produce a smaller embedding. We begin by choosing x_1 to be the point of A satisfying $E_A(x) > x$ for which the support of μ_x extends furthest to the right, and y_1 similarly the point satisfying $E_A(y) < y$ for which the support of μ_y extends furthest to the left. If the support of these measures overlap we show we can exchange mass between μ_{x_1} and μ_{y_1} and embed to find a smaller stopping time. Otherwise we look at those points for which $E_A(x) = x$ and the support overlaps that of μ_{x_1} but extends further to the right. In this way we can find a sequence whose supports overlap (since (15) does not hold) and we may again perform a suitable exchange of mass to show that we can find a smaller embedding. Then we take $x_r = y_1$ and the points satisfy $x_2 < x_3 < \dots < x_{r-1}$.

There are several technical issues we need to address. Firstly, if we find at some stage there are two points which both satisfy the criterion — for example their supports have the same upper bound — then we may use either point. Secondly, if the support of all suitable points has a maximum which is not attained we may still use the same procedure but we must (and can) choose a point which approximates the bound suitably closely for subsequent steps to work. Finally we note that once we choose x_2 , since there is at most one point to the right of y_1 , there exists only a finite number of points left to choose from (by assumption on A) and so the sequence will be finite.

The technical construction is as follows: let x_1 be the largest value such that $E_A(x_1) > x_1$ and

$$\sup\{z : z \in \text{supp}(\mu_{x_1})\} = \sup_{w: E_A(w) > w} \{\sup\{z : z \in \text{supp}(\mu_w)\}\},$$

(or at least so that the left hand side approximates the right hand side sufficiently closely for the next step to work — since the support of the points to the right overlaps we shall be able to find x_1 with supremum of its support sufficiently close to the term on the left) and let y_1 be the smallest value such that $E_A(y_1) < y_1$ and

$$\inf\{z : z \in \text{supp}(\mu_{y_1})\} = \inf_{w: E_A(w) < w} \{\inf\{z : z \in \text{supp}(\mu_w)\}\}.$$

Then (by the assumption that (15) does not hold) we can find a sequence $x_1, x_2, \dots, x_r$ such that $x_r = y_1$ and $x_2 < x_3 < \dots < x_{r-1}$, $E_A(x_i) = x_i$ for $1 < i < r$ and, if we define $I_i = \inf\{\text{intervals } I : \text{supp}(\mu_{x_i}) \subseteq I\}$, then

$$\begin{aligned} \text{Leb}(I_i \cap I_{i+1}) &> 0 & k = 1, \dots, r-1, \\ \text{Leb}(I_i \cap I_{i+2}) &= 0 & k = 1, \dots, r-2. \end{aligned} \tag{16}$$

This is done by choosing at each step the w with $E_A(w) = w$ which overlaps the support of the previous μ_{x_i} and whose support extends furthest to the right, until the support overlaps with the support of μ_{y_1} .

We write $\mu_i = \mu_{x_i}$. For general $1 \leq i < r$ now consider μ'_i defined by

$$\mu'_i = \mu_i|_{(-\infty, y_i)} + \mu_{i+1}|_{(-\infty, y_i)}$$

where y_i is chosen such that $\mu_i([y_i, \infty)) = \mu_{i+1}((-\infty, y_i))$. Then it must be true that $\int w \mu'_i(dw) < \int w \mu_i(dw)$. Define

$$\begin{aligned} m_i &= \int w \mu_i(dw) - \int w \mu'_i(dw) > 0 \\ m_0 &= \int w \mu_1(dw) - \mu_1(\mathbb{R})x_1 > 0 \\ m_r &= \mu_r(\mathbb{R})x_r - \int w \mu_r(dw) > 0 \end{aligned}$$

and set $\Delta m = \inf\{m_i : 0 \leq i \leq r\}$. Then for each i we can find $v_i < z_i$ such that $\mu_i([z_i, \infty)) = \mu_{i+1}((-\infty, v_i))$ and for

$$\mu'_i = \mu_i|_{(-\infty, z_i)} + \mu_{i+1}|_{(-\infty, v_i)}$$

we have

$$\int w \mu_i(dw) - \int w \mu'_i(dw) = \Delta m.$$

Set

$$\begin{aligned} \mu''_1 &= \mu_1|_{(-\infty, z_1)} + \mu_2|_{(-\infty, v_1)}, \\ \mu''_i &= \mu_{i-1}|_{[z_{i-1}, \infty)} + \mu_i|_{[v_{i-1}, z_i)} + \mu_{i+1}|_{(-\infty, v_i)} \quad i = 2, \dots, r-1, \\ \mu''_r &= \mu_{r-1}|_{[z_{r-1}, \infty)} + \mu_r|_{[v_{r-1}, \infty)}. \end{aligned}$$

Then

$$\begin{aligned} \int x \mu''_1 &\geq \mu''_1(\mathbb{R})x_1 \\ \int x \mu''_i &= \mu''_i(\mathbb{R})x_i \quad i = 2, \dots, r-1 \\ \int x \mu''_r &\leq \mu''_r(\mathbb{R})x_r. \end{aligned}$$

So the conditions of Lemma 3.7 are satisfied (due to (16)) for each μ''_i and we can find strictly smaller stopping times on each of the sets $\{T > R(A), R(A) = x_i\}$.

It only remains to show the final statement of the lemma. Let $A' \supset A$ be another set satisfying the conditions of the lemma for some M' , such that $A' \setminus A \subseteq [x, y]$. Then there exists $x', y' \in A'$ such that $x \leq x' < y' \leq y$, $E_{A'}(x') > x'$ and $E_{A'}(y') < y'$ — if this were not the case at least one of the embeddings conditional on $\{R(A') = z\}$ would not be minimal.

Now consider a sequence $A \subset A_1 \subset A_2 \subset \dots$ and such that $A_n \setminus A \subseteq [x, y]$ and $d(z, A_n) \leq 2^{-n}$ for $z \in [x, y]$. Let

$$\begin{aligned} \Lambda &= \{a \in [x, y] : T \leq H_a \text{ on } \{T \geq R(A)\}\}; \\ \Lambda_n &= \{a \in [x, y] : T \leq H_a \text{ on } \{T \geq R(A_n)\}\}. \end{aligned}$$

Then the sets Λ, Λ_n are closed, $\Lambda \supseteq \Lambda_1 \supseteq \Lambda_2 \supseteq \dots$, and each Λ_n is non-empty. So there exists $a_\infty \in \Lambda_n$ for all n . Hence $T \leq H_{a_\infty}$ on $\{T \geq R(A_n)\}$ for all n . But $R(A_n) \downarrow 0$ on $\{B_0 \in [x, y]\}$ and $R(A) \leq H_{a_\infty}$ on $\{B_0 \notin [x, y]\}$. $\square$

This result, although technical in nature, can be thought of as beginning to describe the sort of behaviour we shall expect from minimal embeddings in this general context. The cases considered in Theorem 3.1 suggest behaviour of the form: ‘the process always drifts in the same direction’, if indeed it drifts at all. The example of Remark 3.5 suggests that this is not always possible in the general case, and the previous result suggests that this is modified by breaking the space into two sections, in each of which the process can be viewed separately. The way these sections are determined is clearly dependent on the starting and target measures, and we shall see in the next section that the potential of these measures provides an important tool in determining how this occurs.

4 Minimality and Potential

The main aim of this section is to find equivalent conditions to minimality which allow us to characterise minimality simply in terms of properties of the process $B_{t \wedge T}$. This is partly in order to prove the following result:

The Chacon-Walsh type embedding is minimal when constructed using the functions u_{μ_0} and $c(x) = u_{\mu}(x) - C$ where

$$C = \sup_x \{u_{\mu}(x) - u_{\mu_0}(x)\}. \quad (17)$$

We have already shown that provided the means of our starting and target distribution match, and (9) holds (so that $C = 0$ — the solution in this case to (17)), then the process constructed using the Chacon-Walsh technique is uniformly integrable, and therefore minimal. Of course the Chacon-Walsh construction is simply an example of an embedding, and the functions u_{μ_0} and c are properties solely of the general problem — it seems reasonable however that these functions will appear in the general problem of classifying all minimal embeddings.

So consider a pair μ_0, μ of integrable measures. Remark 2.1(iv) tells us we can choose $C < \infty$ such that (17) holds. We know $u_{\mu_0}(x) - c(x)$ is bounded above, and $\inf_{x \in \mathbb{R}} u_{\mu_0}(x) - c(x) = 0$. We consider

$$\mathcal{A} = \{x \in [-\infty, \infty] : \lim_{y \rightarrow x} [u_{\mu_0}(y) - c(y)] = 0\}. \quad (18)$$

Since both functions are Lebesgue almost-everywhere differentiable, Remark 2.1(v) implies $\mathcal{A} \subseteq \mathcal{A}'$ where $\mathcal{A}'$ is the set

$$\{x \in [-\infty, \infty] : \mu((-\infty, x)) \leq \mu_0((-\infty, x)) \leq \mu_0((-\infty, x]) \leq \mu((-\infty, x])\}. \quad (19)$$

One consequence of this is that if the starting distribution has an atom at a point of $\mathcal{A}$ then the target distribution has an atom at least as large. Also we introduce the following definition. Given a measure ν , $a \in \mathbb{R}$ and $\theta \in [\nu((-\infty, a)), \nu((-\infty, a])]$ we define the measure

$\check{\nu}^{a,\theta}$ to be the measure which is ν on $(-\infty, a)$, has support on $(-\infty, a]$ and $\check{\nu}^{a,\theta}(\mathbb{R}) = \theta$. We also define $\hat{\nu}^{a,\theta} = \nu - \check{\nu}^{a,\theta}$. Then for $a \in \mathcal{A}$ we may find θ such that

$$\begin{aligned}\check{\mu}^{a,\theta}((-\infty, a]) &= \check{\mu}_0^{a,\theta}((-\infty, a]) \\ \hat{\mu}^{a,\theta}([a, \infty)) &= \hat{\mu}_0^{a,\theta}([a, \infty)).\end{aligned}$$

When $\mu_0((-\infty, a)) < \mu_0((-\infty, a])$ there will exist multiple θ . We will occasionally drop the θ from the notation since this is often unnecessary.

These definitions allows us to write the potential in terms of the new measures (for any suitable θ)

$$u_\mu(x) = \int_{(-\infty, x]} (y - x) \check{\mu}^x(dy) + \int_{[x, \infty)} (x - y) \hat{\mu}^x(dy). \quad (20)$$

As a consequence of this and a similar relation for u_{μ_0} , we are able to deduce the following important facts about the set $\mathcal{A}$:

- if $x < z$ are both elements of $\mathcal{A}$ (possibly $\pm\infty$), then

$$\int y (\mu - \check{\mu}^{x,\theta} - \hat{\mu}^{z,\phi})(dy) = \int y (\mu_0 - \check{\mu}_0^{x,\theta} - \hat{\mu}_0^{z,\phi})(dy). \quad (21)$$

That is, we may find measures agreeing with μ and μ_0 on (x, z) and with support on $[x, z]$ which have the same mean.

- If $x \in \mathcal{A}$, by definition

$$u_\mu(x) - u_{\mu_0}(x) \geq \lim_{z \rightarrow -\infty} (u_\mu(z) - u_{\mu_0}(z)). \quad (22)$$

This can be rearranged, using (20), to deduce

$$\int_{(-\infty, x]} y \check{\mu}_0^x(dy) \leq \int_{(-\infty, x]} y \check{\mu}^x(dy)$$

with equality if and only if there is also equality in (22) — that is when $-\infty \in \mathcal{A}$.

Together these imply that the set $\mathcal{A}$ divides $\mathbb{R}$ into intervals on which the starting and target measures place the same amount of mass. Further, the means of the distributions agree on these intervals except for the first (resp. last) interval where the mean of the target distribution will be larger (resp. smaller) than that of the starting distribution unless $-\infty$ (resp. ∞) is in $\mathcal{A}$, when again they will agree. Note the connection between this idea and Lemma 3.8

Before we prove the result we establish several results that are needed in the proof.

Proposition 4.1. *Suppose $T \leq H_{a_\infty}$ is an embedding of μ for $a_\infty \in \mathbb{R}$. Then $a_\infty \in \mathcal{A}$.*

Proof. Clearly a_∞ must lie in $\mathcal{A}'$ (see (19)). Suppose also that $a_\infty < z \in \mathcal{A}$. We may choose θ, ϕ such that $\mu_0 - \check{\mu}_0^{a_\infty, \theta} - \hat{\mu}_0^{z, \phi}$ has no atom at either a_∞ or z .

Then

$$u_{\mu_0}(a_\infty) \geq u_\mu(a_\infty) - C \quad (23)$$

and $C = u_\mu(z) - u_{\mu_0}(z)$ imply

$$\int y (\mu - \check{\mu}^{a_\infty, \theta} - \hat{\mu}^{z, \phi})(dy) \geq \int y (\mu_0 - \check{\mu}_0^{a_\infty, \theta} - \hat{\mu}_0^{z, \phi})(dy), \quad (24)$$

the term on the right being equal to $\mathbb{E}(B_0; B_0 \in (a_\infty, z))$ and the term on the left at most $\mathbb{E}(B_T; B_0 \in (a_\infty, z))$. However $B_T = B_{T \wedge H_{a_\infty}}$ is a supermartingale on $\{B_0 \geq a_\infty\}$, so we must have equality in (24) and hence in (23). So $a_\infty \in \mathcal{A}$. A similar proof can be used for $z < a_\infty$. If there does not exist any such z , $a_\infty \in \mathcal{A}$, since $\mathcal{A} \neq \emptyset$. $\square$

Proposition 4.2. *Suppose T is minimal and A is a countable subset of $\mathbb{R}$ such that A has finitely many elements in every compact subset of $\mathbb{R}$ and $d(x, A) < M$ for all $x \in \mathbb{R}$ and some $M > 0$. Suppose also that $S \leq T$ is a stopping time and $I \subseteq \mathbb{R}$ is an interval such that $\partial I \subseteq A$. If*

$$\mathbb{E}(B_T; F \cap \{B_0 \in I\}) > \mathbb{E}(B_S; F \cap \{B_0 \in I\}) \quad (25)$$

for some $F \in \mathcal{F}_S$ then $E_A(x) > x$ for some $x \in A \cap \bar{I}$.

Proof. We may assume $F \subseteq \{B_0 \in I\}$ and we note that therefore $B_{R(A)} \in A \cap \bar{I}$ on $\{R(A) < T\} \cap F$. Since $B_{t \wedge R(A)}$ is uniformly integrable,

$$\begin{aligned} \mathbb{E}(B_S; F) &= \mathbb{E}(B_{R(A)}; F \cap \{S \leq R(A)\}) + \mathbb{E}(B_S; F \cap \{R(A) < S\}) \\ \mathbb{E}(B_T; F) &= \mathbb{E}(B_{R(A)}; F \cap \{T = R(A)\}) + \mathbb{E}(B_T; F \cap \{R(A) < T\}). \end{aligned}$$

So (25) and the above identities imply

$$\begin{aligned} \mathbb{E}(B_T; F \cap \{R(A) < T\}) &> \mathbb{E}(B_{R(A)}; F \cap \{S \leq R(A) < T\}) \\ &+ \mathbb{E}(B_S; F \cap \{R(A) < S\}). \end{aligned}$$

However if $E_A(x) \leq x$ for all $x \in A \cap \bar{I}$ and T is minimal, by Theorem 3.1:

$$\begin{aligned} \mathbb{E}(B_T; F \cap \{R(A) < S\}) &\leq \mathbb{E}(B_S; F \cap \{R(A) < S\}) \\ \mathbb{E}(B_T; F \cap \{S \leq R(A) < T\}) &\leq \mathbb{E}(B_{R(A)}; F \cap \{S \leq R(A) < T\}) \end{aligned}$$

and we deduce a contradiction. $\square$

Proposition 4.3. *Suppose $F \in \mathcal{F}_0$, $\mathbb{E}(B_T; F) = \mathbb{E}(B_0; F)$ and*

$$\mathbb{E}(B_T | \mathcal{F}_S) \leq B_S \text{ on } F \quad (26)$$

for all stopping times S . Then in fact we have equality — that is

$$\mathbb{E}(B_T | \mathcal{F}_S) = B_S$$

almost surely on F .

Proof. If $\mathbb{P}(F) = 0$ there is nothing to prove. Otherwise we may condition on F to reduce to showing the result when $F = \Omega$.

By the upward martingale theorem (Rogers and Williams, 2000)[Theorem II.69.5], the process

$$X_t = \mathbb{E}(B_T | \mathcal{F}_t)$$

is uniformly integrable. Also $\mathbb{E}(B_T | \mathcal{F}_0) \leq B_0$ and $\mathbb{E}(B_T) = \mathbb{E}(B_0)$ implies $\mathbb{E}(B_T | \mathcal{F}_0) = B_0$. Let $Y_t = B_{T \wedge t} - X_{T \wedge t}$. By (26) Y_t is a non-negative local martingale such that $Y_0 = Y_T = 0$. Hence $Y \equiv 0$. $\square$

Lemma 4.4. *If T is minimal and $a \in \mathcal{A}$ then $T \leq H_a$ and*

$$\mathbb{E}(B_T | \mathcal{F}_S) \leq B_S \text{ on } \{B_0 \geq a\}; \quad (27)$$

$$\mathbb{E}(B_T | \mathcal{F}_S) \geq B_S \text{ on } \{B_0 \leq a\}. \quad (28)$$

Proof. Suppose initially $a \in \mathbb{R}$. Let $\theta = \mu_0((-\infty, a))$. If $\{B_0 < a\} \not\subseteq \{B_T \leq a\}$ a.s. then also $\{B_0 \geq a\} \not\subseteq \{B_T \geq a\}$ a.s. and

$$\begin{aligned} \mathbb{E}(B_0; B_0 < a) &= \int y \check{\mu}_0^{a, \theta}(dy) \leq \int y \check{\mu}^{a, \theta}(dy) < \mathbb{E}(B_T; B_0 < a); \\ \mathbb{E}(B_0; B_0 \geq a) &= \int y \hat{\mu}_0^{a, \theta}(dy) \geq \int y \check{\mu}^{a, \theta}(dy) > \mathbb{E}(B_T; B_0 < a). \end{aligned}$$

So there exists $x_1 \leq a$ and $x_2 \geq a$ such that (by Proposition 4.2)

$$E_A(x_1) < x_1 \text{ and } E_A(x_2) > x_2$$

for a suitable choice of A — a contradiction to Lemma 3.8.

A similar argument can be used with $\theta = \mu_0((-\infty, a])$ to deduce that $\{B_0 \leq a\} \subseteq \{B_T \leq a\}$ a.s. and $\{B_0 \geq a\} \subseteq \{B_T \geq a\}$ a.s.. So if there is an atom of μ_0 at a then paths starting at a must also stop at a , and hence (by the minimality of T) must stop immediately — i.e. $T = 0$ on $\{B_0 = a\}$.

So consider paths for which $\{B_0 < a\}$. For almost all these paths, for some choice of A , $B_{R(A)} < a$. If (28) fails, by Proposition 4.2 there exists $x < a$ such that $E_A(x) < x$. Then Lemma 3.8 and (for $\theta = \mu_0((-\infty, a))$)

$$\int y \check{\mu}_0^{a, \theta}(dy) \leq \int y \check{\mu}^{a, \theta}(dy)$$

imply there must also exist $y < x$ such that $E_A(y) > y$, and hence $a' < a$ such that $T \leq H_{a'}$. Then $B_{t \wedge T}$ is a supermartingale on $\{B_0 > a'\}$ (and a submartingale on $\{B_0 \leq a'\}$). But Proposition 4.1 and (21) imply $\mathbb{E}(B_0; a' < B_0 < a) = \mathbb{E}(B_T; a' < B_0 < a)$ and therefore (by Proposition 4.3) $B_{t \wedge T}$ is a true martingale on $\{a' < B_0 < a\}$ — in particular $T \leq H_a$ on $\{B_0 < a\}$, and (28) holds. Similarly (27) can be shown to hold.

So suppose now that $a = \infty$ (the case $a = -\infty$ is similar) and there exists $a' < \infty$ also in $\mathcal{A}$. By the above, $T \leq H_{a'}$ and so $B_{t \wedge T}$ is a supermartingale on $\{B_0 > a'\}$, while by (21) $\mathbb{E}(B_0; B_0 > a') = \mathbb{E}(B_T; B_0 > a')$, and hence $B_{t \wedge T}$ satisfies (28) by Proposition 4.3.

Finally suppose $\mathcal{A} = \{\infty\}$. By Lemma 3.8 $E_A(x) \geq x$ for all suitable choices of A and all x . Hence, by Proposition 4.2,

$$\mathbb{E}(B_T|\mathcal{F}_S) \geq B_S.$$

□

We note that some of the above arguments, particularly the use of Proposition 4.3, allow us to deduce that if there exists $a \in \mathcal{A}$, $|a| < \infty$ for which $T \leq H_a$ then (27) and (28) hold and $T \leq H_{a'}$ for all $a' \in \mathcal{A}$.

Lemma 4.5. *Suppose that for all stopping times S with $S \leq T$ and $\mathbb{E}|B_S| < \infty$ we have*

$$\mathbb{E}(B_T|\mathcal{F}_S) \leq B_S \quad a.s.. \quad (29)$$

Then T is minimal.

We refer the reader to Lemma 8 of Cox and Hobson (2003), the proof of which is still valid in the more general case.

Of course we may replace the ' $\leq$ ' in (29) with ' $\geq$ ' or ' $=$ ' without altering the conclusion.

Lemma 4.6. *If $T \leq H_{\mathcal{A}} = \inf\{t \geq 0 : B_t \in \mathcal{A}\}$ is a stopping time of the Brownian motion $(B_t)_{t \geq 0}$ where $B_0 \sim \mu_0$ and $B_T \sim \mu$, and*

$$\mathbb{E}(B_T|\mathcal{F}_S) \leq B_S \quad \text{on } \{B_0 \geq a_-\} \quad (30)$$

$$\mathbb{E}(B_T|\mathcal{F}_S) \geq B_S \quad \text{on } \{B_0 \leq a_+\}, \quad (31)$$

where $a_- = \inf \mathcal{A}$ and $a_+ = \sup \mathcal{A}$, then T is minimal.

Proof. Choose $a \in \mathcal{A}$. By assumption $T \leq H_a$ and by Lemma 4.5 T is minimal for $\check{\mu}^a$ on $\{B_0 \leq a\}$ and for $\hat{\mu}^a$ on $\{B_0 \geq a\}$. It must then be minimal for μ . □

These results show the equivalence of minimality and the conditions in (30), (31). The following theorem states this together with some extra equivalent conditions. It should be thought of as the extension of Theorem 3.1 to the setting with a general starting measure.

Theorem 4.7. *Let B be a Brownian motion such that $B_0 \sim \mu_0$ and T a stopping time such that $B_T \sim \mu$, where μ_0, μ are integrable. Let $\mathcal{A}$ be the set defined in (18) and $a_+ = \sup\{x \in [-\infty, \infty] : x \in \mathcal{A}\}$, $a_- = \inf\{x \in [-\infty, \infty] : x \in \mathcal{A}\}$. Then the following are equivalent:*

(i) T is minimal;

(ii) $T \leq H_{\mathcal{A}}$ and for all stopping times $R \leq S \leq T$

$$\mathbb{E}(B_S|\mathcal{F}_R) \leq B_R \quad \text{on } \{B_0 \geq a_-\}$$

$$\mathbb{E}(B_S|\mathcal{F}_R) \geq B_R \quad \text{on } \{B_0 \leq a_+\};$$

(iii) $T \leq H_{\mathcal{A}}$ and for all stopping times $S \leq T$

$$\begin{aligned}\mathbb{E}(B_T | \mathcal{F}_S) &\leq B_S \text{ on } \{B_0 \geq a_-\} \\ \mathbb{E}(B_T | \mathcal{F}_S) &\geq B_S \text{ on } \{B_0 \leq a_+\};\end{aligned}$$

(iv) $T \leq H_{\mathcal{A}}$ and for all $\gamma > 0$

$$\begin{aligned}\mathbb{E}(B_T; T > H_{-\gamma}, B_0 \geq a_-) &\leq -\gamma \mathbb{P}(T > H_{-\gamma}, B_0 \geq a_-) \\ \mathbb{E}(B_T; T > H_{\gamma}, B_0 \leq a_+) &\geq \gamma \mathbb{P}(T > H_{\gamma}, B_0 \leq a_+);\end{aligned}$$

(v) $T \leq H_{\mathcal{A}}$ and as $\gamma \rightarrow \infty$

$$\begin{aligned}\gamma \mathbb{P}(T > H_{-\gamma}, B_0 \geq a_-) &\rightarrow 0 \\ \gamma \mathbb{P}(T > H_{\gamma}, B_0 \leq a_+) &\rightarrow 0.\end{aligned}$$

We begin by proving the following result:

Proposition 4.8. *If (v) holds and $S \leq T$ then $\mathbb{E}|B_S| < \infty$.*

Proof. We show that $\mathbb{E}(|B_S|; B_0 \geq a_-) < \infty$. Since $B_{t \wedge H_{-k}}$ is a supermartingale on $\{B_0 \geq -k\}$,

$$\begin{aligned}\mathbb{E}(B_{T \wedge H_{-k}}; B_S < 0, S < H_{-k}, B_0 \geq a_- \wedge (-k)) \\ \leq \mathbb{E}(B_{S \wedge H_{-k}}; B_S < 0, S < H_{-k}, B_0 \geq a_- \wedge (-k)).\end{aligned}$$

The term on the left hand side is equal to:

$$\begin{aligned}\mathbb{E}(B_T; B_S < 0, T < H_{-k}, B_0 \geq a_- \wedge (-k)) \\ - k \mathbb{P}(B_S < 0, S \leq H_{-k} < T, B_0 \geq a_- \wedge (-k)).\end{aligned}$$

The first term converges (by dominated convergence) to $\mathbb{E}(B_T; B_S < 0, B_0 \geq a_-)$ and the second term vanishes by the assumption. By monotone convergence

$$\begin{aligned}\mathbb{E}(B_S; B_S < 0, B_0 \geq a_-) &= \lim_k \mathbb{E}(B_S; B_S < 0, S < H_{-k}, B_0 \geq a_- \wedge (-k)) \\ &\geq \lim_k \mathbb{E}(B_T; B_S < 0, S < H_{-k}, B_0 \geq a_- \wedge (-k)) \\ &\geq \mathbb{E}(B_T; B_S < 0, B_0 \geq a_-) \geq -\mathbb{E}(B_T^-) > -\infty.\end{aligned}$$

Also

$$\begin{aligned}\mathbb{E}(B_0; B_0 \geq a_- \wedge (-k)) &\geq \mathbb{E}(B_{S \wedge H_{-k}}; B_0 \geq a_- \wedge (-k)) \\ &= \mathbb{E}(B_S; B_0 \geq a_- \wedge (-k), S < H_{-k}) \\ &\quad - k \mathbb{P}(H_{-k} \leq S, B_0 \geq a_- \wedge (-k)),\end{aligned}$$

and

$$\begin{aligned}\mathbb{E}(B_S; B_0 \geq a_- \wedge (-k), S < H_{-k}) &= \mathbb{E}(B_S^+; B_0 \geq a_- \wedge (-k), S < H_{-k}) \\ &\quad - \mathbb{E}(B_S^-; B_0 \geq a_- \wedge (-k), S < H_{-k}),\end{aligned}$$

so

$$\begin{aligned}\mathbb{E}(B_S^+; B_0 \geq a_- \wedge (-k), S < H_{-k}) &\leq \mathbb{E}(B_0; B_0 \geq a_- \wedge (-k)) \\ &\quad + \mathbb{E}(B_S^-; B_0 \geq a_- \wedge (-k), S < H_{-k}) \\ &\quad + k\mathbb{P}(H_{-k} \leq S, B_0 \geq a_- \wedge (-k)).\end{aligned}$$

By monotone and dominated convergence, in the limit we have

$$\begin{aligned}\mathbb{E}(B_S^+; B_0 \geq a_-) &\leq \mathbb{E}(B_0; B_0 \geq a_-) + \mathbb{E}(B_S^-; B_0 \geq a_-) \\ &< \infty.\end{aligned}$$

So $\mathbb{E}(|B_S|; B_0 \geq a_-) < \infty$. Similarly $\mathbb{E}(|B_S|; B_0 \leq a_+) < \infty$, and together these imply $\mathbb{E}(B_S) < \infty$. $\square$

Proof of Theorem 4.7. Clearly $(ii) \implies (iii) \implies (iv) \implies (v)$ (the final implication following from dominated convergence). We also know $(i) \iff (iii)$. We show $(v) \implies (ii)$.

Suppose $A \in \mathcal{F}_R$, $A \subseteq \{B_0 \geq a_-\}$ and set $A_k = A \cap \{R < H_{-k}\} \cap \{B_0 \geq -k\}$. Then

$$\mathbb{E}(B_{S \wedge H_{-k}}; A_k) \leq \mathbb{E}(B_{R \wedge H_{-k}}; A_k).$$

By Proposition 4.8 we may apply dominated convergence to deduce that in the limit as $k \rightarrow \infty$ the right-hand side converges to $\mathbb{E}(B_R; A)$. Also

$$\begin{aligned}\mathbb{E}(B_{S \wedge H_{-k}}; A_k) &= \mathbb{E}(B_S; A \cap \{B_0 \geq -k\} \cap \{S \leq H_{-k}\}) \\ &\quad + k\mathbb{P}(A, R < H_{-k} < S, B_0 \geq -k),\end{aligned}$$

where the second term converges to zero by assumption and the first converges to $\mathbb{E}(B_S; A)$ by dominated convergence. $\square$

5 Minimality of the Limit

We will want to show that stopping times constructed using the techniques of Section 2 are indeed minimal when (17) is satisfied. To deduce that a stopping time T constructed using the balayage techniques is minimal, we approximate T by the sequence of stopping times T_n given in the construction (so T_1 is the exit time from the first interval we construct, and so on). Then it is clear that the stopping times T_n satisfy the conditions of Lemma 4.6, since they are either the first exit time from a bounded interval, or the first time to leave $(-\infty, \alpha]$ (resp. $[\beta, \infty)$) for some $\alpha < a_-$ (resp. $\beta > a_+$). Our aim is then to deduce that the limit is minimal.

Proposition 5.1. *Suppose that T_n embeds μ_n , μ_n converges weakly to μ and $\mathbb{P}(|T_n - T| > \varepsilon) \rightarrow 0$ for all $\varepsilon > 0$. Then T embeds μ .*

If also $l_n \rightarrow l_\infty < \infty$ where $l_n = \int |x| \mu_n(dx)$ and $l_\infty = \int |x| \mu(dx)$, and T_n is minimal for μ_n , then T is minimal for μ .

Remark 5.2. Since $\mu_n \implies \mu$, on some probability space we are able to find random variables X_n and X with laws μ_n and μ such that $X_n \rightarrow X$ a.s.. By Scheffé's Lemma therefore

$$\mathbb{E}|X_n - X| \rightarrow 0 \text{ if and only if } \mathbb{E}|X_n| \rightarrow \mathbb{E}|X|,$$

the second statement being equivalent to $l_n \rightarrow l_\infty$ in the statement of Proposition 5.1.

Before we prove this result, we will show a useful result on the distribution of the maximum. This will be used in the proof of Proposition 5.1, and also be important for the work in the next section, when we will show that the inequality in (32) can be attained by a class of stopping times created by balayage techniques.

Lemma 5.3. *Let T be a minimal embedding of μ in a Brownian motion started with distribution μ_0 . Then for all $x \in \mathbb{R}$*

$$\mathbb{P}(\bar{B}_T \geq x) \leq \inf_{\lambda < x} \frac{1}{2} \left[1 + \frac{u_{\mu_0}(x) - c(\lambda)}{x - \lambda} \right]. \quad (32)$$

Proof. Define the stopping time $\bar{H}_x = \inf\{t \geq 0 : B_t \geq x\}$, the first time that B_t goes above x . Then we note the following inequality, which (by considering on a case by case basis) holds for all paths and all pairs $\lambda < x$:

$$\mathbf{1}_{\{\bar{B}_T \geq x\}} \leq \frac{1}{x - \lambda} \left[B_{T \wedge \bar{H}_x} + \frac{|B_T - \lambda| - (B_T + \lambda)}{2} - \frac{|B_0 - x| + (B_0 - x)}{2} \right]. \quad (33)$$

In particular, on $\{\bar{B}_T < x\}$, when therefore $\{B_0 < x\}$:

$$0 \leq \frac{1}{x - \lambda} \left[B_T + \begin{cases} -\lambda & : B_T > \lambda \\ -B_T & : B_T \leq \lambda \end{cases} \right]. \quad (34)$$

While on $\{\bar{B}_T \geq x\}$,

$$\begin{aligned} 1 &\leq \frac{1}{x - \lambda} \left[B_{T \wedge \bar{H}_x} + \begin{cases} -\lambda & : B_T > \lambda \\ -B_T & : B_T \leq \lambda \end{cases} - \begin{cases} B_0 - x & : B_0 > x \\ 0 & : B_0 \leq x \end{cases} \right] \\ &\leq \frac{1}{x - \lambda} \left[x + \begin{cases} -\lambda & : B_T > \lambda \\ -B_T & : B_T \leq \lambda \end{cases} \right]. \end{aligned} \quad (35)$$

So we may take expectations in (33) to get

$$\mathbb{P}(\bar{B}_T \geq x) \leq \frac{1}{2} \left[1 + \frac{2\mathbb{E}(B_{T \wedge \bar{H}_x}) + (u_{\mu_0}(x) - u_\mu(\lambda)) - (\mathbb{E}(B_T) + \mathbb{E}(B_0))}{(x - \lambda)} \right]. \quad (36)$$

We can deduce (32) provided we can show

$$C \geq 2\mathbb{E}(B_{T \wedge \bar{H}_x}) - (\mathbb{E}(B_T) + \mathbb{E}(B_0)) \quad (37)$$

since (36) holds for all $\lambda < x$.

We now consider $a \in \mathcal{A}$ possibly taking the values $\pm\infty$. Since $u_\mu(a) - u_{\mu_0}(a) = C$ for $a \in \mathcal{A}$, we can deduce

$$C = 2\mathbb{E}(B_T; B_T \geq a) + 2\mathbb{E}(B_0; B_0 < a) - \mathbb{E}(B_T) - \mathbb{E}(B_0)$$

where we note that $\{B_T < a\} = \{B_0 < a\}$. Theorem 4.7 tells us that

$$\mathbb{E}(B_{T \wedge \bar{H}_x}; B_0 < a) \leq \mathbb{E}(B_T; B_0 < a) \quad (38)$$

$$\mathbb{E}(B_{T \wedge \bar{H}_x}; B_0 \geq a) \leq \mathbb{E}(B_0; B_0 \geq a) \quad (39)$$

and (37) holds. $\square$

We also have the following result:

Proposition 5.4. *Suppose μ and $\{\mu_n\}_{n \geq 1}$ are all integrable distributions such that $\mu_n \Rightarrow \mu$ and $l_n = \int |y| \mu_n(dy) \rightarrow \int |y| \mu(dy) = l_\infty$. Then u_{μ_n} converges uniformly to u_μ .*

Proof. Fix $\varepsilon > 0$. By (20), using the fact that $\mu - \hat{\mu} = \check{\mu}$ we may write

$$u_\mu(x) = \int_{-\infty}^{\infty} (x - y) \mu(dy) + 2 \int_{-\infty}^x (y - x) \mu(dy) = x - \int_{-\infty}^{\infty} y \mu(dy) + 2 \int_{-\infty}^x (y - x) \mu(dy),$$

and similarly for u_{μ_n} , hence

$$u_{\mu_n}(x) - u_\mu(x) = (m_\infty - m_n) + 2 \int_{-\infty}^x (y - x) (\mu_n - \mu)(dy), \quad (40)$$

where we write m_n, m_∞ for the means of μ_n and μ respectively; $m_n \rightarrow m_\infty$ as a consequence of Remark 5.2. Since μ is integrable, as $x \downarrow -\infty$,

$$\int_{-\infty}^x (x - y) \mu(dy) \downarrow 0.$$

By (40) and Lemma 2.2 (which implies u_{μ_n} converges to u_μ pointwise, the C in (4) being 0 since $l_n \rightarrow l_\infty$), for all $x \in \mathbb{R}$

$$\int_{-\infty}^x (x - y) \mu_n(dy) \rightarrow \int_{-\infty}^x (x - y) \mu(dy)$$

as $n \rightarrow \infty$. Finally we note that both sides of the above are increasing in x .

Consider

$$|u_{\mu_n}(x) - u_\mu(x)| \leq |m_\infty - m_n| + 2 \int_{-\infty}^x (x - y) \mu_n(dy) + 2 \int_{-\infty}^x (x - y) \mu(dy).$$

We may choose x_0 sufficiently small that $\int_{-\infty}^{x_0} (x_0 - y) \mu(dy) < \varepsilon$, and therefore such that

$$\int_{-\infty}^x (x - y) \mu(dy) \leq \int_{-\infty}^{x_0} (x_0 - y) \mu(dy) < \varepsilon$$

for all $x \leq x_0$. By the above and Remark 5.2 we may now choose $n_0(\varepsilon)$ such that for all $n \geq n_0(\varepsilon)$

$$|m_\infty - m_n| < \varepsilon \text{ and } \left| \int_{-\infty}^{x_0} (x_0 - y) \mu_n(dy) - \int_{-\infty}^{x_0} (x_0 - y) \mu(dy) \right| < \varepsilon.$$

Then for all $x \leq x_0$ and for all $n \geq n_0(\varepsilon)$,

$$|u_{\mu_n}(x) - u_\mu(x)| \leq \varepsilon + 2 \times 2\varepsilon + 2\varepsilon = 7\varepsilon.$$

Similarly we can find $x_1, n_1(\varepsilon)$ such that $|u_{\mu_n}(x) - u_\mu(x)| \leq 7\varepsilon$ for all $x \geq x_1$ and all $n \geq n_1(\varepsilon)$. Finally u_{μ_n}, u_μ are both Lipschitz and pointwise $u_{\mu_n}(x) \rightarrow u_\mu(x)$ and we must have uniform convergence on any bounded interval, and in particular on $[x_0, x_1]$. $\square$

Proof of Proposition 5.1. Suppose first that there exists $a \in \mathcal{A} \cap \mathbb{R}$. We show that $T \leq H_a$ for all such a . As usual, we write μ_0 for the starting measure, and $c(x) = u_\mu(x) - C$. We define C_n to be the smallest value such that $u_{\mu_0}(x) \geq u_{\mu_n}(x) - C_n$ and the functions $c_n(x) = u_{\mu_n}(x) - C_n$. Note that $l_n = u_{\mu_n}(0)$, so $\lim_{n \rightarrow \infty} u_{\mu_n}(0)$ exists. Then (by Lemma 2.2(i) or equivalently (Chacon, 1977)[Lemma 2.5]) weak convergence implies

$$\lim_{n \rightarrow \infty} u_{\mu_n}(x) = u_\mu(x) - K$$

for all $x \in \mathbb{R}$ and (here) $K = 0$ since $u_{\mu_n}(0) \rightarrow u_\mu(0)$.

By Lemma 5.3 for $x \in \mathbb{R}$ and $\lambda < x$

$$\mathbb{P}(\overline{B}_{T_n} \geq x) \leq \frac{1}{2} \left[1 + \frac{u_{\mu_0}(x) - u_{\mu_n}(\lambda) + C}{x - \lambda} + \frac{C_n - C}{x - \lambda} \right],$$

and we take the limit as $n \rightarrow \infty$, using Proposition 5.4 (so that $C_n \rightarrow C$) and noting that $\mathbb{P}(\overline{B}_{T_n} \geq x) \rightarrow \mathbb{P}(\overline{B}_T \geq x)$, to get

$$\mathbb{P}(\overline{B}_T \geq x) \leq \frac{1}{2} \left[1 + \frac{u_{\mu_0}(x) - c(\lambda)}{x - \lambda} \right].$$

Suppose now $x = a$. Since the above holds for all $\lambda < a$, we may take the limit of the right hand side as $\lambda \uparrow a$, in which case $u_{\mu_0}(a) = c(a)$, and by Remark 2.1(v)

$$\begin{aligned} \mathbb{P}(\overline{B}_T \geq a) &\leq \frac{1}{2} [1 + c'_-(a)] \\ &\leq \frac{1}{2} [1 + (1 - 2\mu((-\infty, a)))] \\ &\leq \mu([a, \infty)). \end{aligned}$$

By considering $-B_t$ we may deduce that $\mathbb{P}(\underline{B}_T \leq a) \leq \mu((-\infty, a])$. Hence $\mathbb{P}(T \leq H_a) = 1$, and we deduce that T is minimal.

It only remains to show (by Lemma 4.5) that if $\infty \in \mathcal{A}$ then

$$\mathbb{E}(B_T | \mathcal{F}_S) \geq B_S$$

for all stopping times $S \leq T$. The case where $-\infty \in \mathcal{A}$ follows from $B_t \mapsto -B_t$. In particular, for $S \leq T$ and $A \in \mathcal{F}_S$ we need to show

$$\mathbb{E}(B_T; A) \geq \mathbb{E}(B_S; A). \quad (41)$$

In fact we need only show the above for sets $A \subseteq \{S < T\}$ since it clearly holds on $\{S = T\}$. So we can define $A_n = A \cap \{S < T_n\}$ and therefore $\mathbb{P}(A \setminus A_n) \rightarrow 0$ as $n \rightarrow \infty$. Also $A_n \in \mathcal{F}_{S \wedge T_n}$. By Theorem 4.7 and the fact that the T_n are minimal

$$\begin{aligned} \mathbb{E}(B_{S \wedge T_n}; A_n) &\leq \mathbb{E}(B_{T_n}; A_n \cap \{B_0 \leq a_+^n\}) + \mathbb{E}(B_{S \wedge T_n}; B_0 > a_+^n) \\ &\quad - \mathbb{E}(B_{S \wedge T_n}; A_n^C \cap \{B_0 > a_+^n\}) \\ &\leq \mathbb{E}(B_{T_n}; A_n \cap \{B_0 \leq a_+^n\}) \mathbb{E}(B_0; B_0 > a_+^n) \\ &\quad - \mathbb{E}(B_{T_n}; A_n^C \cap \{B_0 > a_+^n\}) \\ &\leq \mathbb{E}(B_{T_n}; A_n) - \mathbb{E}(B_{T_n}; \{B_0 > a_+^n\}) + \mathbb{E}(B_0; B_0 > a_+^n) \end{aligned}$$

where a_+^n is the supremum of the set $\mathcal{A}_n$ (that is the corresponding set to $\mathcal{A}$ for the measures μ_0, μ_n). This is not necessarily infinite.

So it is sufficient for us to show that

$$\lim_n \mathbb{E}(B_{T_n}; A_n) = \mathbb{E}(B_T; A); \quad (42)$$

$$\lim_n \mathbb{E}(B_S; A_n) = \mathbb{E}(B_S; A), \quad (43)$$

and

$$\lim_n |\mathbb{E}(B_0; B_0 > a_+^n) - \mathbb{E}(B_{T_n}; B_0 > a_+^n)| = 0. \quad (44)$$

For (42) we consider $|\mathbb{E}(B_T; A) - \mathbb{E}(B_{T_n}; A_n)|$. Then

$$|\mathbb{E}(B_T; A) - \mathbb{E}(B_{T_n}; A_n)| \leq \mathbb{E}(|B_T|; A \setminus A_n) + \mathbb{E}(|B_T - B_{T_n}|; A_n)$$

and the first term tends to zero by dominated convergence (this follows from the assumption that T_n converges to T in probability). For the second term we show $\mathbb{E}(|B_T - B_{T_n}|) \rightarrow 0$. Fix $\varepsilon > 0$. We have

$$|B_T - B_{T_n}| \leq |B_{T_n}| - |B_T| + 2|B_T| \mathbf{1}_{\{|T_n - T| \geq \varepsilon\}} + 2|B_T - B_{T_n}| \mathbf{1}_{\{|T_n - T| \leq \varepsilon\}}.$$

We take expectations and let $n \rightarrow \infty$. By the definition of μ_n the first two terms cancel each other out, while the third tends to zero by dominated convergence. For the last term, by the (strong) Markov property

$$\mathbb{E}(|B_T - B_{T_n}|; |T_n - T| \leq \varepsilon) \leq \mathbb{E}(|B_\varepsilon|) = \sqrt{\frac{\varepsilon}{2\pi}}.$$

Consequently, in the limit, $\mathbb{E}(|B_T - B_{T_n}|; |T_n - T| \leq \varepsilon) \rightarrow 0$ and (42) holds. We want to apply Lemma 4.5 so we can assume that $\mathbb{E}|B_S| < \infty$, and (43) follows by dominated convergence.

Finally we consider (44). Let $\theta_n = \mu_0((-\infty, a_+^n])$. Since $a_+^n \in \mathcal{A}_n$ we have

$$\begin{aligned}
\mathbb{E}(B_0; B_0 > a_+^n) - \mathbb{E}(B_{T_n}; B_0 > a_+^n) &= \int y \hat{\mu}_0^{a_+^n, \theta_n}(dy) - \int y \hat{\mu}_n^{a_+^n, \theta_n}(dy) \\
&= \int (y - a_+^n) \hat{\mu}_0^{a_+^n, \theta_n}(dy) - \int (y - a_+^n) \hat{\mu}_n^{a_+^n, \theta_n}(dy) \\
&= \frac{1}{2} \left[\int y (\mu_0 - \mu_n)(dy) + u_{\mu_n}(a_+^n) - u_{\mu_0}(a_+^n) \right] \\
&= \frac{1}{2} \left[\int y (\mu_0 - \mu)(dy) - C_n \right],
\end{aligned}$$

where we have used the fact that (for a general measure ν)

$$\int (y - x) \hat{\nu}^x(dy) = \frac{1}{2} \left[\int y \nu(dy) - u_\nu(x) - x \right].$$

As $n \rightarrow \infty$, since $\infty \in \mathcal{A}$,

$$\int y (\mu_0 - \mu_n)(dy) \rightarrow \int y (\mu_0 - \mu)(dy) = C.$$

So we need only show that $C_n \rightarrow C$, which follows from the uniform convergence of u_{μ_n} to u_μ (Proposition 5.4). $\square$

6 Tangents and Azema-Yor Type Embeddings

One of the motivations for this paper is to discuss generalisations of the Azema-Yor family of embeddings (see Azéma and Yor (1979); Jacka (1988)) to the integrable starting/target measures we have discussed already.

The aim is therefore to find the embedding which maximises the law of the maximum, $\sup_{0 \leq t \leq T} B_t$ (or in the more general case $\sup_{0 \leq t \leq T} |B_t|$). If we look for the maximum within the class of all embeddings there is no natural maximum embedding. For this reason we consider the class of minimal embeddings. Lemma 5.3 establishes that there is some natural limit when we consider this restriction. In fact the extended Azema-Yor embedding will attain the limit in (32).

The idea is to use the machinery from the previous sections to show the embeddings exist as limits of the Chacon-Walsh type embeddings of Section 2. It is then possible to show that the embeddings are minimal and that they attain equality in (32).

Theorem 6.1. *If T is a stopping time as described in Lemma 2.6, where C as described in the lemma is*

$$C = \inf_x \{u_\mu(x) - u_{\mu_0}(x)\}, \quad (45)$$

then T is minimal.

Proof. Lemma 2.6 suggests a sequence T_n of stopping times for which T is the limit. We note that we can modify the definition of T_n so that T'_n is specified by the functions

$f_1, f_2, \dots, f_n, f^{-1}, f^{+1}$ without altering their limit (as a consequence of (6)), where f^{-1} is the tangent to g with gradient -1 and f^{+1} is the tangent to g with gradient 1 . It is easy to see that this ensures that $\mathbb{E}(B_{T'_n}) = \mathbb{E}(B_T)$ (by (3)), and also that $u_{\mu_n}(0) \rightarrow u_\mu(0)$ and $n \rightarrow \infty$. Consequently the stopping times T'_n and T satisfy the conditions of Proposition 5.1, where it is clear that the T'_n are all minimal, since each step clearly satisfies the conditions of Theorem 4.7 as a consequence of (45). So T is minimal. $\square$

Define the function

$$\Phi(x) = \operatorname{argmin}_{\lambda < x} \left\{ \frac{u_{\mu_0}(x) - c(\lambda)}{x - \lambda} \right\}. \quad (46)$$

In the cases described by Azéma and Yor (1979), this is the barycentre function. It can also be seen to agree with the function appearing in the generalisation of the Azema-Yor stopping time to non-centred means which appears in Cox and Hobson (2003). A similar function is used in Hobson (1998) who examines the case where starting and target means are centred and satisfy (9). $\Phi(\cdot)$ can be thought of graphically as the point (below x) at which there exists a tangent to $c(\cdot)$ meeting the function $u_{\mu_0}(\cdot)$ at x .

Lemma 6.2. *The Azema-Yor stopping time*

$$T = \inf\{t \geq 0 : B_t \leq \Phi(\overline{B}_t)\} \quad (47)$$

is minimal and attains equality in (32).

We prove this lemma using an extension of an idea first suggested in Meilijson (1983). We approximate T by taking tangents to c , starting with gradient -1 , and increasing to $+1$. As the number of tangents we take increases, the stopping time converges to T . The general approximation sequence can be seen in Figure 1.

Proof. We apply Lemma 2.6 for each n to the functions $f_1^n, f_2^n, \dots, f_{m(n)}^n$, which are chosen as tangents to $c(\cdot)$ with increasing gradients, so that f_1^n has gradient -1 , f_m^n has gradient 1 , and so that the difference in the gradients of consequential tangents is less than $\frac{1}{n}$. We also choose the tangents in such a way that the points at which successive tangents intersect each other (which are B_{T_n} stops) are at most $\frac{1}{n}$ apart when they lie within $[-n, n]$ (at least as far as this is possible — if both μ_0 and μ have an interval containing no mass, it might not be possible to manage this, but this case will not be important). This defines a (minimal) stopping time T_n such that (by (3)) $\mathbb{E}(B_{T_n}) = \int x \mu(dx)$. Also, by considering $\mu_n = \mathcal{L}(B_{T_n})$, $|\mu_n((-\infty, x)) - \mu((-\infty, x))| \leq \frac{1}{n}$ for all $x \in \mathbb{R}$. So $\mu_n \Rightarrow \mu$. The choice of T_n also ensures that $\mathbb{P}(|T - T_n| > \varepsilon) \rightarrow 0$ for all $\varepsilon > 0$. Consequently T is minimal.

To deduce that T attains equality in (32) we note that $\Phi(x)$ is the optimal choice for λ in (32), and by the definition of $\Phi(x)$,

$$\begin{aligned} \{\overline{B}_T < x\} &\subseteq \{B_T \leq \Phi(x)\} \\ \{\overline{B}_T \geq x\} &\subseteq \{B_T \geq \Phi(x)\}. \end{aligned}$$




Figure 1: Approximating the Azema-Yor stopping time: we take tangents to the potential from left to right. In the limit the tangents become closer. The dotted lines highlight the points at which the approximated stopping time will stop the process.

This means we attain equality in (34) and (35), and so only need show that we have equality in (38) and (39) for equality in (32) to hold. But for x given, we may calculate the potential of $\mu' = \mathcal{L}(B_{T \wedge \bar{H}_x})$ — where $\bar{H}_x = \inf\{t \geq 0 : B \geq x\}$ — as:

$$u_{\mu'}(y) = \begin{cases} u_{\mu}(y) & : y \leq \Phi(x); \\ u_{\mu}(\Phi(x)) + \frac{y - \Phi(x)}{x - \Phi(x)}(u_{\mu_0}(x) - u_{\mu}(\Phi(x))) & : \Phi(x) \leq y \leq x; \\ u_{\mu_0}(y) & : y \geq x. \end{cases}$$

It then follows from Theorem 4.7 and (3) that equality holds. $\square$

Remark 6.3. The embedding due to Jacka (1988) can be viewed easily in this framework. Essentially the embedding can be described as follows. We wish to find an embedding which maximises $\mathbb{P}(\sup_{t \leq T} |B_t| \geq x)$ simultaneously for all x , subject to T being minimal. The optimal construction can be described thus in the Chacon-Walsh picture: choose the tangent to u_{μ} with gradient 0. Let (a, b) be the points at which the tangent intersects $c(x)$, where C is chosen to be the value given by (17); the first step of the construction is to run the process until it leaves the interval $I = (a, b)$. The construction will now have at least two separate halves; on the positive half we run the Azema-Yor construction of (47), and on the negative half we run the reverse of the Azema-Yor construction. Such a construction will therefore be minimal, and can be shown to be optimal — for details we refer the reader to Cox and Hobson (2003).




Figure 2: The Chacon-Walsh type picture for an approximation to the Vallois stopping time, with a general starting distribution. We note that after the first two steps, there could still be mass at the extremes. This mass will have to be embedded using some suitable procedure — for example a Vallois construction using the local time at a different level.

This technique can be trivially extended to maximising $\mathbb{P}(\sup_{t \leq T} f(B_t) \geq x)$, where f is a function which is increasing above some point x_0 and decreasing below x_0 .

Remark 6.4. In a slightly different vein, the construction of Vallois (1983) with decreasing functions can also be seen in this framework. Choose $\varepsilon > 0$ and construct alternate tangents; tangents of positive gradient intersecting the current potential at ε , and tangents of negative gradient intersecting the current potential at 0 (see Figure 2). This can be repeated a number of times, and results in an approximation of $c(x)$; suitable further choices of tangents can be used to construct a full embedding. If c touches u_{μ_0} at finite points, any choice of construction outside the interval containing 0 may be used. By construction this is a minimal embedding.

The Vallois construction results on taking the limit as $\varepsilon \downarrow 0$; the appearance of the local time in the construction results since the limiting embeddings are determined by the number of downcrossings of $[0, \varepsilon]$, which has as a limit the local time (see Revuz and Yor (1999)[VI.1.10]). An application of Proposition 5.1 allows us to deduce that this limit is minimal. For further details we refer the reader to Cox (2004).

References

- Azéma, J. and M. Yor (1979), Une solution simple au problème de Skorokhod, in ‘Séminaire de Probabilités, XIII (Univ. Strasbourg, Strasbourg, 1977/78)’, Vol. 721 of *Lecture Notes in Math.*, Springer, Berlin, pp. 90–115.
- Bertoin, J. and Y. Le Jan (1992), ‘Representation of measures by balayage from a regular recurrent point’, *Ann. Probab.* **20**(1), 538–548.
- Chacon, R. V. (1977), ‘Potential processes’, *Trans. Amer. Math. Soc.* **226**, 39–58.
- Chacon, R. V. and J. B. Walsh (1976), One-dimensional potential embedding, in ‘Séminaire de Probabilités, X (Prèmiere partie, Univ. Strasbourg, Strasbourg, année universitaire 1974/1975)’, Springer, Berlin, pp. 19–23. *Lecture Notes in Math.*, Vol. 511.
- Cox, A. M. G. (2004), Skorokhod embeddings: non-centred target distributions, diffusions and minimality, PhD thesis, University of Bath. Available from <http://www-users.york.ac.uk/~amgc500>.
- Cox, A. M. G. and D. G. Hobson (2003), Skorokhod embeddings, minimality and non-centred target distributions. University of Bath Mathematics Preprints 03/11, <http://www.maths.bath.ac.uk/MATHEMATICS/preprints.html>.
- Dubins, L. E. (1968), ‘On a theorem of Skorohod’, *Ann. Math. Statist.* **39**, 2094–2097.
- Hobson, D. G. (1998), The maximum maximum of a martingale, in ‘Séminaire de Probabilités, XXXII’, Vol. 1686 of *Lecture Notes in Math.*, Springer, Berlin, pp. 250–263.
- Jacka, S. D. (1988), ‘Doob’s inequalities revisited: a maximal H^1 -embedding’, *Stochastic Process. Appl.* **29**(2), 281–290.
- Meilijson, I. (1983), On the Azéma-Yor stopping time, in ‘Seminar on probability, XVII’, Vol. 986 of *Lecture Notes in Math.*, Springer, Berlin, pp. 225–226.
- Monroe, I. (1972), ‘On embedding right continuous martingales in Brownian motion’, *Ann. Math. Statist.* **43**, 1293–1311.
- Oblój, J. (2004), The Skorokhod embedding problem and its offspring. Preprint [arxiv.org:math.PR/0401114](http://arxiv.org/math.PR/0401114).
- Perkins, E. (1986), The Cereteli-Davis solution to the H^1 -embedding problem and an optimal embedding in Brownian motion, in ‘Seminar on stochastic processes, 1985 (Gainesville, Fla., 1985)’, Birkhäuser Boston, Boston, MA, pp. 172–223.
- Revuz, D. and M. Yor (1999), *Continuous martingales and Brownian motion*, Vol. 293 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*, third edn, Springer-Verlag, Berlin.

- Rogers, L. C. G. and D. Williams (2000), *Diffusions, Markov processes, and martingales. Vol. 1*, Cambridge University Press, Cambridge. Foundations, Reprint of the second (1994) edition.
- Root, D. H. (1969), ‘The existence of certain stopping times on Brownian motion’, *Ann. Math. Statist.* **40**, 715–718.
- Skorokhod, A. V. (1965), *Studies in the theory of random processes*, Translated from the Russian by Scripta Technica, Inc, Addison-Wesley Publishing Co., Inc., Reading, Mass.
- Vallois, P. (1983), Le problème de Skorokhod sur $\mathbf{R}$: une approche avec le temps local, in ‘Seminar on probability, XVII’, Vol. 986 of *Lecture Notes in Math.*, Springer, Berlin, pp. 227–239.
- Vallois, P. (1992), ‘Quelques inégalités avec le temps local en zero du mouvement brownien’, *Stochastic Process. Appl.* **41**(1), 117–155.